

Guidance for the interpretation of elevated serum potassium in primary care

Patients at risk of TRUE hyperkalaemia or those at greater risk of its effects:

- CKD /Deterioration in renal function from previous
- Relevant drugs (see box below)
- Diabetes
- Metabolic acidosis
- Patients with acute illness e.g. AKI
- Older patients
- Patients with a cardiac history

Relevant Drugs

- **K retaining drugs:** ACEi, ARB, Spironolactone, NSAIDS, Amiloride and other K sparing diuretics; anti-fungals
- **K supplement:** Lo-salt
- **K containing drugs:** Movicol, Fybogel

Consider SPURIOUS HYPERKALAEMIA if:

- Prolonged tourniquet application/fist clenching
 - Traumatic venepuncture
 - Delayed separation exacerbated by low temperature
 - Sample stored in fridge before dispatch
 - Possible EDTA contamination – consider if low Calcium/Magnesium/ALP
 - Young, well patient
 - Isolated, marked elevation of potassium with normal/no change in renal function
 - Normal acid/base (bicarbonate normal)
 - No relevant drugs (see left column)
 - Elevated white cells or platelets
- Note:** Potassium results from **haemolysed** samples are not reported.

Potassium results ≥ 6.5 mmol/L will be phoned as a critical result