

Classification: Official

Publication approval reference: PAR1621_ix

National shared care protocol:

Azathioprine and mercaptopurine for patients within adult services (non-transplant indications)

4 July 2022, Version 1

Review date – January 2025

The content of this shared care protocol was correct as of January 2022. As well these protocols, please ensure that [summaries of product characteristics](#) (SPCs), [British national formulary](#) (BNF) or the [Medicines and Healthcare products Regulatory Agency](#) (MHRA) or [NICE](#) websites are reviewed for up-to-date information on any medicine.

Specialist responsibilities

- Assess the patient and provide diagnosis; ensure that this diagnosis is within scope of this shared care protocol ([section 2](#)) and communicated to primary care.
- Use a shared decision making approach; discuss the benefits and risks of the treatment with the patient and provide the appropriate counselling (see [section 11](#)) to enable the patient to reach an informed decision. Obtain and document patient consent. Provide an appropriate patient information leaflet.
- Assess for contraindications and cautions (see [section 4](#)) and interactions (see [section 7](#)).
- Conduct required baseline investigations and initial monitoring (see [section 8](#)).
- Initiate and optimise treatment as outlined in [section 5](#).
- Transfer to primary care is normally after the patient has been treated for 3 months and with satisfactory investigation results for at least 4 weeks. Prescribe sufficient medication to enable transfer to primary care, including where there are unforeseen delays to transfer of care.
- Once treatment is optimised, complete the shared care documentation and send to patient's GP practice detailing the diagnosis, current and ongoing dose, baseline and most recent test results, confirm the monitoring schedule and when the next monitoring is required. Include contact information ([section 13](#)).
- Conduct the required monitoring in [section 8](#) and communicate the results to primary care. After each review, advise primary care whether treatment should be continued, confirm the ongoing dose, and whether the ongoing monitoring outlined in [section 9](#) remains appropriate.
- Give advice to primary care on continuing treatment if a woman becomes or wishes to become pregnant or breastfeed.
- Provide advice to primary care on the management of adverse effects if required.

Primary care responsibilities

- Respond to the request from the specialist for shared care in writing. It is asked that this be undertaken within 14 days of the request being made, where possible.
- If accepted, prescribe ongoing treatment as detailed in the specialists request and as per [section 5](#) taking into any account potential drug interactions in [section 7](#).
- Adjust the dose of azathioprine or mercaptopurine prescribed as advised by the specialist.
- Conduct the required monitoring as outlined in [section 9](#).
- Assess for possible interactions with azathioprine or mercaptopurine when starting new medicines (see [section 7](#)).
- Manage any adverse effects as detailed in [section 10](#) and discuss with specialist team when required.
- Stop azathioprine or mercaptopurine and discuss urgently with the specialist if bone marrow suppression is suspected.
- Discuss other adverse effects with the specialist team as clinically appropriate (see [section 10](#)).
- Contact the specialist team for advice if the patient becomes or plans to become pregnant.
- Stop treatment as advised by the specialist.

Patient and/or carer responsibilities

- Take azathioprine or mercaptopurine as prescribed and do not stop taking it without speaking to their primary care prescriber or specialist.
- Tell anyone who prescribes them a medicine that they are taking azathioprine or mercaptopurine.
- Attend regularly for monitoring and review appointments with primary care and specialist. Be aware that medicines may be stopped if they do not attend appointments.
- Report adverse effects to their primary care prescriber. Seek immediate medical attention if they develop any symptoms as detailed in [section 11](#).
- Report the use of any over the counter medications to their prescriber and be aware they should discuss the use of azathioprine or mercaptopurine with their pharmacist before purchasing any OTC medicines.

Inform the specialist or primary care prescriber as soon as possible if they become pregnant or wish to become pregnant.

1. Background

[Back to top](#)

Azathioprine and mercaptopurine are disease modifying anti-rheumatic drugs (DMARDs). They are used as immunosuppressant anti-metabolites either alone or, more commonly, in combination with other agents (usually corticosteroids) to influence the immune response. Therapeutic effect may be evident only after weeks or months and can include a steroid sparing effect, thereby reducing the toxicity associated with high dosage and prolonged usage of corticosteroids.

Azathioprine and mercaptopurine are not licensed for all the conditions they are used to treat, as noted below. However, their use for the indications below are established and supported by various sources and bodies including the BNF, NICE, British Society for Rheumatology (BSR) and British Health Professionals in Rheumatology (BHPR), British Association of

Dermatologists (BAD), British Thoracic Society (BTS), Association of British Neurologists (ABN) and British Society of Gastroenterology (BSG).

2. Indications

[Back to top](#)

Azathioprine

The licensed indications for azathioprine include:

- Auto-immune chronic active hepatitis
- Auto-immune haemolytic anaemia
- Chronic refractory idiopathic thrombocytopenic purpura
- Dermatomyositis
- Inflammatory bowel disease (IBD)
- Pemphigus vulgaris
- Polyarteritis nodosa
- Polymyositis
- Pyoderma gangrenosum
- Rheumatoid arthritis
- Systemic lupus erythematosus (SLE)

Licensed indications vary with brand. See relevant summary of product characteristics ([see SPC](#)) for full details.

This shared care protocol also includes treatment of chronic inflammatory conditions where off-label use of azathioprine is appropriate, including, but not limited to the following specialities and conditions:

- Dermatology (e.g. severe eczema)
- Neurology (e.g. myasthenia gravis, demyelinating conditions)
- Ophthalmology (e.g. uveitis, scleritis)
- Oral medicine (e.g. Behçet's disease, refractory inflammatory oral disease)
- Renal medicine (e.g. immune-mediated nephritis)
- Respiratory disease (e.g. interstitial lung disease)
- Rheumatology (e.g. inflammatory arthritis, connective tissue disease, vasculitis, giant cell arteritis)

These indications are off-label. The initiating specialist must specify the indication for each patient when initiating shared care and clearly state when use is off-label.

Mercaptopurine

This shared care protocol includes treatment of chronic inflammatory conditions where off-label use of mercaptopurine is appropriate, including, but not limited to the following conditions:

- Inflammatory bowel disease

- Autoimmune encephalitides
- Autoimmune hepatitis

These indications are off-label. The specialist must specify the indication for each patient when initiating shared care and clearly state when use is off-label.

This shared care protocol applies to adults aged 18 and over. It does not include use of azathioprine or mercaptopurine for transplant or oncology indications.

3. **Locally agreed off-label use**

[Back to top](#)

To be agreed and completed locally (include supporting information)

4. **Contraindications and cautions**

[Back to top](#)

This information does not replace the Summary of Product Characteristics (SPC), and should be read in conjunction with it. Please see [BNF](#) & [SPC](#) for comprehensive information.

Contraindications:

- Known hypersensitivity to the active substance or any excipients. Hypersensitivity to 6-mercaptopurine (6-MP) should alert the prescriber to probable hypersensitivity to azathioprine.
- Absent or very low thiopurine methyltransferase (TPMT) activity – risk of life-threatening pancytopenia.

Cautions:

- Live vaccines (e.g. oral polio, oral typhoid, MMR, BCG, yellow fever): should be avoided in patients taking azathioprine at a dose greater than 3 mg/kg/day, or mercaptopurine greater than 1.5 mg/kg/day. Please refer to the [Green Book Chapter 6](#) for current advice regarding the use of live vaccines in patients taking immune modulators. Contact the specialist if further guidance is required.
- Patients with active/history of pancreatitis.
- Concomitant prescribing of allopurinol: A 75% dose reduction of azathioprine/mercaptopurine is required, see [section 7](#).
- Patients receiving azathioprine or mercaptopurine are at an increased risk of developing lymphoproliferative disorders and other malignancies, notably skin cancers, sarcomas and uterine cervical cancer in situ. Exposure to sunlight and UV light should be limited and patients should wear protective clothing and use a sunscreen with a high protection factor to minimize the risk of skin cancer and photosensitivity
- Patients with low thiopurine methyltransferase (TPMT) activity are at increased risk of myelosuppression. Substantial dose reduction is generally required.
- Severe infection.

- Severely impaired hepatic or bone marrow function.
- Pregnancy and breastfeeding (see [section 12](#)).

Treatment may need to be monitored more frequently in the following:

- Elderly patients
- Impaired renal function
- Mild/moderately impaired hepatic function
- Mild/moderately impaired bone marrow function

5. Initiation and ongoing dose regimen

[Back to top](#)

- Transfer of monitoring and prescribing to primary care is normally after at least 12 weeks, and when the patient's dose has been optimised and with satisfactory investigation results for at least 4 weeks.
- The duration of treatment & frequency of review will be determined by the specialist, based on clinical response and tolerability.
- All dose or formulation adjustments will be the responsibility of the initiating specialist unless directions have been discussed and agreed with the primary care clinician.
- Termination of treatment will be the responsibility of the specialist.

There is a wide dose range depending on the indication. The selected dose will be tailored to the individual patient and decided by the specialist.

The initial stabilisation period must be prescribed by the initiating specialist.

Transfer of monitoring and prescribing to primary care is usually after 3 months. The duration of treatment will be determined by the specialist based on clinical response and tolerability.

Maintenance dose (following initial stabilisation):

Usual dose range:

- **Azathioprine: 0.5–3 mg/kg daily**, adjusted according to response.
- **Mercaptopurine: 1-1.5mg/kg daily**, adjusted according to response.

Some patients may respond to lower doses. Please note patients may be initiated on more than one DMARD.

The initial maintenance dose must be prescribed by the initiating specialist.

Conditions requiring dose adjustment:

Lower doses may be required if there is significant renal or hepatic impairment, in elderly patients, and in patients with mild/moderately impaired bone marrow function, TPMT deficiency or NUDT15 mutation ([see SPC](#)).

6. Pharmaceutical aspects

[Back to top](#)

Route of administration:	Oral
Formulation:	Azathioprine 25mg and 50mg tablets Azathioprine 10 mg/mL oral suspension (Jayempi®) Mercaptopurine 50mg tablets Mercaptopurine 20mg/ml oral suspension (Xaluprine®) Mercaptopurine 10mg tablets (unlicensed import)
Administration details:	The tablets should be swallowed whole and not split / crushed. Can be taken either with or without food, but patients should standardise which method is chosen. Tablets should be taken at least 1 hour before or 2 hours after milk or dairy products. Taking with or after food may relieve nausea, however the oral absorption of azathioprine or mercaptopurine may be reduced. Consideration should be given to monitoring therapeutic efficacy more closely if patient is taking azathioprine or mercaptopurine consistently with food. For azathioprine or mercaptopurine oral suspension, the bottle should be shaken vigorously for at least 30 seconds to ensure the suspension is well mixed.
Other important information:	Providing the film coating of azathioprine tablets remains intact, there is no risk or additional precautions required when handling tablets. Azathioprine and mercaptopurine are cytotoxic. It is recommended that they are handled following local recommendations for the handling and disposal of cytotoxic agents. Mercaptopurine tablets and oral suspension are not bioequivalent with respect to peak plasma concentration; increased haematological monitoring is advised if switching between formulations. When prescribing mercaptopurine, remain vigilant with regards to the similarity in name with mercaptamine.

7. Significant medicine interactions

[Back to top](#)

The following list is not exhaustive. Please see [BNF](#) or [SPC](#) for comprehensive information and recommended management.

The following drugs must not be prescribed without consultation with the specialist:

- Allopurinol** has the potential to cause thiopurine toxicity and should be avoided, except with specialist input. Allopurinol may be recommended in combination with thiopurines by the specialist for IBD patients, particularly in those who are unable to tolerate to or do not respond to treatment with a thiopurine alone. The dose of

azathioprine or mercaptopurine should be reduced by 75% if used concurrently with allopurinol. If considering prescribing allopurinol, discuss with the specialist for advice and a dose adjustment.

- **Febuxostat** has the potential to cause thiopurine toxicity; avoid in combination with azathioprine or mercaptopurine.
- **Live vaccines** (e.g. oral polio, oral typhoid, MMR, BCG, yellow fever) can be given to patients on stable long term low dose corticosteroid therapy (defined as $\leq 20\text{mg}$ prednisolone per day for >14 days) alone or in combination with low dose non-biological oral immune modulating drugs (e.g. azathioprine up to 3mg/kg/day or mercaptopurine up to 1.5mg/kg/day). Clinician discretion is advised. Please refer to the [Green Book Chapter 6](#) for current advice, and advice for patients taking higher doses.
- **Warfarin** – thiopurines may reduce anticoagulant effects of warfarin.
- **Co-trimoxazole / trimethoprim** – possible increased risk of haematological toxicity, however evidence is conflicting and this combination is often used in practice.
- **Clozapine** - avoid due to increased risk of agranulocytosis.
- **Ribavirin** - increased risk of haematological toxicity when azathioprine given concurrently and this combination should be avoided.
- **Aminosalicylates** (sulfasalazine, mesalazine or olsalazine) - increased risk of haematological toxicity with concomitant thiopurine due to TPMT inhibition. Dose adjustment of azathioprine or mercaptopurine and additional monitoring of FBC may be required.

The following drugs may be prescribed with caution:

- **ACE inhibitors** - increase the risk of anaemia and or leukopenia.

Cimetidine and indomethacin - concomitant administration of thiopurines may increase the risk of myelosuppression.

8. Baseline investigations, initial monitoring and ongoing monitoring to be undertaken by specialist

[Back to top](#)

Monitoring at baseline and during initiation is the responsibility of the specialist; only once the patient is optimised on the chosen medication with no anticipated further changes expected in immediate future will prescribing and monitoring be transferred to primary care.

Baseline investigations:

- Height and weight
- Blood pressure
- Full blood count (FBC)
- Urea and electrolytes (U&Es) & creatinine clearance (CrCl)
- Alanine aminotransferase (ALT) and/or aspartate aminotransferase (AST), and albumin
- Baseline thiopurine methyl transferase (TPMT) status
- Screening for viral infections as per local policy, e.g. HIV, hepatitis B and C, varicella zoster, Epstein Barr virus, cytomegalovirus

- Screening for lung disease, including tuberculosis, should be undertaken at clinician discretion on a case by case basis
- Confirm cervical screening is up to date

Provide or request appropriate vaccination prior to treatment initiation, according to local arrangements (e.g. pneumococcal, shingles, influenza, COVID-19)

Initial monitoring and at dose change:

To be repeated every 2 weeks until the dose has been stable for 6 weeks, then monthly for 3 months:

- FBC
- U&Es, including creatinine and CrCl
- LFTs, including AST and/or ALT, and albumin

Following a dose increase repeat every 2 weeks until the dose has been stable for 6 weeks, then revert to previous schedule.

More frequent monitoring is appropriate in patients at higher risk of toxicity.

Ongoing monitoring:

The specialist will retain the responsibility for monitoring the patient's ongoing response to treatment and advise if a dose change or treatment cessation is appropriate. This should usually be undertaken annually.

After each review, advise primary care whether treatment should be continued, confirm the ongoing dose, and whether the ongoing monitoring outlined in [section 9](#) remains appropriate.

9. Ongoing monitoring requirements to be undertaken by primary care

[Back to top](#)

See [section 10](#) for further guidance on management of adverse effects/responding to monitoring results.

Monitoring and actions	Frequency
• FBC • U&Es including creatinine and CrCl • ALT and/or AST, and albumin • Rheumatology patients: CRP &/or ESR specialist to confirm (both are not routinely required)	Monthly for three months, unless already completed in secondary care. Thereafter at least every 12 weeks, and more frequently in patients at higher risk of toxicity, as advised by the specialist team. The exact frequency of monitoring to be communicated by the specialist in all cases.
Patients aged 70-79 years old could be eligible for the shingles vaccine (varicella zoster). For patients taking prednisolone exceeding 20mg daily or azathioprine exceeding 3mg/kg/day a non-live vaccine should be used. Specialist input may be required. If patient is taking additional DMARDs, check advice for all drugs. Please refer to Green Book Chapter 6 and Chapter 28a (Shingles) for further details.	• Shingles vaccination: one-off. • Influenza vaccination: annual. It is advisable to add the patient to the influenza vaccine list. COVID-19 vaccination as per national schedule.

Annual influenza (The Green Book, Chapter 19) vaccinations are recommended COVID-19 vaccination is safe and recommended (see The Green Book, Chapter 14a). Repeat pneumococcal vaccine may be indicated. See Green Book Chapter 25 for advice.	
(If relevant) If monitoring results are forwarded to the specialist team, please include clear clinical information on the reason for sending, to inform action to be taken by secondary care.	
10. Adverse effects and other management Back to top Any serious adverse reactions should be reported to the MHRA via the Yellow Card scheme. Visit www.mhra.gov.uk/yellowcard For information on incidence of ADRs see relevant summaries of product characteristics	
Result	Action for primary care
As well as responding to absolute values in laboratory tests, a rapid change or a consistent trend in any value should prompt caution and extra vigilance	
Full blood count: • White blood cells less than $3.5 \times 10^9/L$ • Lymphocytes less than $0.5 \times 10^9/L$ • Neutrophils less than $1.6 \times 10^9/L$ • Platelets less than $140 \times 10^9/L$ Eosinophilia greater than $0.5 \times 10^9/L$	Discuss urgently with specialist team, and consider interruption. NB: Isolated lymphopenia or eosinophilia is often a feature of the underlying autoimmune indication, and is rarely an indication to discontinue azathioprine.
Mean cell volume >105 fl NB: Reversible, dose-related increases in mean corpuscular volume are a known effect of thiopurines.	Consider interruption in treatment if there is a significant increase from baseline. Check serum folate, B12, alcohol history and TSH and treat any underlying abnormality. If results of these additional investigations are normal discuss with specialist team urgently.

Signs or symptoms of bone marrow suppression, e.g. unexplained bleeding or bruising with or without sore throat, mouth ulcers	Consider interruption in treatment. Check FBC immediately and discuss with the specialist team. See haematological monitoring above.
Infections: Infection requiring antibiotics	Temporarily withhold thiopurine until the patient has recovered. Consider additional investigations (e.g. FBC), if clinically appropriate.
Liver function tests: ALT or AST >100 units/L, or any sudden increases (e.g. double of baseline), OR Unexplained fall in serum albumin <30g/L Jaundice	Withhold and discuss with specialist team. When used for hepatology indications, continue treatment and discuss with specialist urgently. Check any other reason for risk of hepatic dysfunction such as alcohol history and drug interactions, including OTC or complementary medication.
Renal function: - Creatinine rise >30% over 12 months, or calculated GFR reduces to <60ml/min	Withhold and discuss with specialist team
Gastrointestinal disorders: Nausea	Review for reversible causes. Advise patient to take with food. If no improvement contact specialist team.
Suspected pancreatitis	Withhold and discuss with specialist team.

11. Advice to patients and carers

[Back to top](#)

The specialist will counsel the patient with regard to the benefits and risks of treatment and will provide the patient with any relevant information and advice, including patient information leaflets on individual medicines.

The patient should be advised to report any of the following signs or symptoms to their primary care prescriber without delay:

Signs or symptoms indicating haematological toxicity, e.g. sore throat, infection, unexplained or abnormal bruising or bleeding.

Signs or symptoms of pancreatitis, e.g. abdominal pain, nausea, or vomiting

Signs of symptoms of hepatic toxicity, e.g. Jaundice (yellowing of the skin or whites of the eyes)

The patient should be advised to:

- During a serious infection azathioprine or mercaptopurine should be temporarily discontinued until the patient has recovered from the infection.
- That vaccination in line with current national advice (e.g. for COVID-19, influenza) is safe and recommended.
- Tell anyone who prescribes them a medicine that they are taking azathioprine or mercaptopurine. Always ask a pharmacist before purchasing any medicines over the counter, including herbal remedies, and ask if they are safe.
- To inform their specialist or primary care prescriber promptly if pregnancy occurs or is planned.
- All women aged 25-64 years old should be encouraged to participate in national cervical cancer screening programmes. There is no need to attend more frequently than recommended.
- Patients have a small increased risk of skin cancers so should be advised to wear high factor sunscreen and to wear a hat and protective clothing when in strong sunshine. Sun beds should be avoided. Patients should be advised to carry out regular self-examination of the skin and report if there are any new lesions and/or changes to skin.
- Patients taking azathioprine at a dose of 3 mg/kg or more, or mercaptopurine at a dose of 1.5 mg/kg/day or more should be advised to avoid contact with people with chicken pox or shingles and report any such contact urgently to their primary care prescriber. If the patient is exposed, contact the specialist for advice. For detailed advice on risk assessment and post exposure prophylaxis following exposure to chicken pox and shingles, see:

the [Green Book \(Chapter 34\)](#)

UKSHA guidance: [Guidelines on post exposure prophylaxis \(PEP\) for varicella/shingles April 2022](#)

Patient information:

- General information: <https://www.nhs.uk/medicines/azathioprine/>
- General information: <https://patient.info/medicine/azathioprine-azapress-imuran>
- Rheumatology: <https://www.versusarthritis.org/about-arthritis/treatments/drugs/azathioprine/>
- Dermatology: <https://www.bad.org.uk/for-the-public/patient-information-leaflets/azathioprine>
- Patient information leaflets are also available from <https://www.medicines.org.uk/emc/search?q=azathioprine>

Gastroenterology:

- <https://www.crohnsandcolitis.org.uk/about-crohns-and-colitis/publications/azathioprine-mercaptopurine>
- <https://gutscharity.org.uk/advice-and-information/conditions/crohns-disease/>
- <https://gutscharity.org.uk/advice-and-information/conditions/ulcerative-colitis/>

12. Pregnancy, paternal exposure and breast feeding

[Back to top](#)

It is the responsibility of the specialist to provide advice on the need for contraception to male and female patients on initiation and at each review, but the ongoing responsibility for providing this advice rests with both the primary care prescriber and the specialist.

All patients should be informed of the risks and benefits of taking this medicine during pregnancy and breastfeeding. The specialist team should be contacted if a patient becomes pregnant or is planning to become pregnant or breastfeed.

the following:

Pregnancy:

The [BSR and BHPR guideline on prescribing DMARDs in pregnancy and breastfeeding](#) advises that azathioprine is compatible throughout pregnancy at doses $\leq 2\text{mg/kg/day}$.

Current available data do not suggest that mercaptopurine exposure during pregnancy increases the risk of miscarriage, congenital malformation, intrauterine death, fetal growth restriction, or preterm delivery but the data are limited for some outcomes. A careful assessment of risk versus benefit should be made before mercaptopurine is prescribed to patients who are pregnant.

The [British Society of Gastroenterology consensus guidelines on the management of inflammatory bowel disease](#) advises that both maintenance and flares can be treated as normal with thiopurines (azathioprine and mercaptopurine) during pregnancy.

Information for healthcare professionals: <https://www.medicinesinpregnancy.org/bumps/monographs/USE-OF-AZATHIOPRINE-OR-MERCAPTOPURINE-IN-PREGNANCY/>

Information for patients and carers: <https://www.medicinesinpregnancy.org/Medicine--pregnancy/Azathioprinemercaptopurine/>

Breastfeeding:

Azathioprine is compatible with breastfeeding, although the active metabolite mercaptopurine is present in breast milk. A risk versus benefit assessment is advised. If used during breastfeeding, monitor for signs of infection or immunosuppression. If high doses of azathioprine are used, monitor infant blood counts. If mercaptopurine is used, monitor infant's blood count and liver function.

Information for healthcare professionals:

- <https://www.sps.nhs.uk/medicines/azathioprine/>
- <https://www.sps.nhs.uk/medicines/mercaptopurine/>

Paternal exposure:

Azathioprine and mercaptopurine are compatible with paternal exposure. There is currently no evidence of adverse fetal effects relating to paternal use.

- Information for healthcare professionals: <https://www.medicinesinpregnancy.org/bumps/monographs/PATERNAL-USE-OF-AZATHIOPRINE-OR-MERCAPTOPURINE/>

13. Specialist contact information

[Back to top](#)

Speciality	Contact name	Contact Details
Haematology	ESNEFT Ipswich Site	
	Dr Debo Ademokun Dr Isobel Chalmers Dr Andrew Hodson Dr Ioana Nitu-Whalley Dr Mahesh Prahladan	Email Secreateries: Haem.Secs@esneft.nhs.uk Team tel: 01473 703718
	ESNEFT Colchester Site	
	Dr Gavin Campbell Dr Michael Hamblin Dr Mahalakshmi Mohan Dr Joseph Padayatty Dr Khalid Saja	Email secretaries: HaematologyPA@esneft.nhs.uk Tel secretaries: 01206 742357
Neurology	ESNEFT Ipswich Site	
	Dr Simon Kerrigan Dr Petr Pokorny Dr Mortada Elyas Dr Clare Galton Dr Andrew Graham	Email Secretaries: neurologyipswich@esneft.nhs.uk Tel: 01473 704044 / 01473 704041
	ESNEFT Colchester Site	
	Dr Imtiaz Ahmed Dr Dr Natalia Salari Dr Mohammed Sharief Epilepsy Nurse Specialist	Telephone Secretaries: 01206 742243 Email Secretaries: ch.neurology@esneft.nhs.uk Tel Epilepsy Nurse Specialist: 01206 746472
Gastroenterology	WSFT	
	Dr Smriti Agarwal Dr Charlotte Brierley Dr Francesca Crawley Dr Alexandre Costa Dr Kate Maresh Dr Edward Needham Dr Dunya Luke	Tel: 01284 712984 Tel: 01284 712984 Tel: 01284 712897 Tel: 01284 712846 Tel: 01284 712984 Tel: 01284 712897 Tel: 01284 712846 Email: NeurologySecretaries@wsh.nhs.uk
Gastroenterology	ESNEFT Ipswich Site	

	Dr Louise Scovell Dr Abdul Mohsen Dr Hermant Laxaman Dr Anne Robins Dr Khalid Yousif Dr Bohdan Semiginovsky IBD Nurse Specialist	Tel: 01473 703519 Tel: 01473 702535 Tel: 01473 703188 Tel: 01473 703505 Tel: 01473 702114 Tel: 01473 702865 Email: gastrosecs@esneft.nhs.uk Advice and guidance on NHS e-referrals
	ESNEFT Colchester Site	
	Dr Ian Gooding Dr Donagh O'Riordan Dr Mary McStay Dr Achuth Shenoy Dr Simon Ralphs IBD Nurse Specialist	Email secretaries: chu- ftr.gastroenterologysecretaries@esneft.nhs.uk 01206 744231 Email: colchesteribdadviceline@esneft.nhs.uk
	ESNEFT Ipswich Site	
	Shuk Yates, Clinical Nurse Specialist Dermatology Jodie Newman, Clinical Nurse Specialist Dermatology Dr Liberta Labinoti	Consultant (via secretaries): 01473 704042 Specialist nurse: 01473 706877 Email: IPS.derm@esneft.nhs.uk Advice and guidance on NHS e-referrals
	WSFT	
	Dr Krishna Basavaraju Dr Miriam Fort Gasia Dr Abdullah Mawas Dr Dan Sharpstone Dr Nadim Sheikh Dr Simon Whalley Dr Banerjee Dr Gavin IBD Specialist Nurses	Tel: 01284 713650 Tel: 01284 712901 Tel: 01284 712901 Tel: 01284 713481 Tel: 01284 713650 Tel: 01284 713461 Tel: 01284 712877 Tel: 01284 713481 Tel: 01284 712611 Email Nurses: IBD-Nurses@wsh.nhs.uk Email Secretaries: GastroenterologySecretaries@wsh.nhs.uk
Dermatology	ESNEFT Colchester Site	
	Dr Farooq Dr Asfar Dr Galahitiyawa, Kerry Penfold, Clinical Nurse Specialist Dermatology Soumya Sunny, Clinical Nurse Specialist Dermatology Dr Khalid Mahmood Susan Robertson, Clinical Nurse Specialist Dermatology	Email: dermatologycolchester@esneft.nhs.uk Specialist Nurse: 01206 487 216

	WSFT	
	Dr Lis Andrews-Fraser Dr Jacqui Hawkins Dr Rachel Jenkins Dr Aruni Ranasinghe Dr Kun Chen Dr Abbas Al-Shammari Nurse Specialist Dermatology	Email: dermatology@wsh.nhs.uk
Renal	ESNEFT Ipswich Site	
	Dr. Brian Camilleri Dr Gerald Glancey Dr. Praveen Ande Dr Richard Smith	Team Tel: 01473 704117 Email Nurses: Renal.Nurses@esneft.nhs.uk
	ESNEFT Colchester Site	
	Dr Gabor Cserep Dr Dimitrios Kirmizis	Email: RenalSecretaries@esneft.nhs.uk Tel: 01206 742612
	WSFT	
	Dr William Petchey Dr Vivian Yiu Dr Hsu Pheen Chong	Team Tel: 01284 713148
Ophthalmology	ESNEFT Ipswich Site	
	Ramy Bassily Richard Goble Vasileios Kostakis Henry Z.H. Lin Kuranageri Poornesh Deepak Vayalambone Dr Sonja Mansfield Smith Paula Varley	Email secretaries: eyesecretaries@esneft.nhs.uk or Paula.Varley@esneft.nhs.uk Team Tel: 01473 702597/703502
	ESNEFT Colchester Site	
	Vivek Bansal Chrysostomos (Chrys) D. Dimitriou Gopan Ghosh Henry Z.H. Lin Achyut Mukherjee Mahmoud Radwan	Email: chu-otr.eyedepartment@esneft.nhs.uk Team Tel: 01206 746860
Rheumatology	ESNEFT Ipswich Site	
	Dr Lane / Dr Watts: Dr Weerakoon / Dr Shand Dr Abdulkader/Dr Ntatsaki	Tel: 01473 702362 Tel: 01473 702039 Tel: 01473 707324 Email: ihn-tr.rheum-sharedcare@nhs.net Advice and guidance on NHS e-referrals
	ESNEFT Colchester Site	

Dr Paul Byrne Dr Tom Walton Dr Rachel Davies Dr Lauren Steel Clinical Nurse Specialist May Clark Clinical Nurse Specialist Lucy Mascurine		Tel Department: (01206) 742165/742279 Email Nurses: rheumatology.nurseadvice@esneft.nhs.uk
ESNEFT Ipswich Site		
Dr Matthew Burton Dr Jonathan Douse Dr Nicholas Innes Dr Rana Rabbani Dr Yu Hon Man Dr Md Erfanur Shuvo		Email Secretaries : respsecretaries@esneft.nhs.uk Tel Secretaries: 01473 703779 / 01473 703508
WSFT		
Dr Katerina Achilleos Dr Catherine Bevington Dr Shweta Bhagat Dr Vivek Rajagopal		Tel: 01284 713484 Tel: 01284 712907 Tel: 01284 712977 Tel: 01284 712590 Email Nurses: RheumatologyNurses2@wsh.nhs.uk Tel: 01284713707 Email Secretary: RheumatologySecretaries@wsh.nhs.uk
Respiratory	ESNEFT Colchester Site	
	Respiratory Dr Rekha Badiger Dr Samantha Cooper Dr Peter Hawkins Dr Misha Sidhu Dr Mphatso Wirima Dr Emma Boynton Dr Safin Rahman	Email Secretaries: respsecretaries@esneft.nhs.uk Tel: 01206 742612
14. Additional information Back to top		
Where patient care is transferred from one specialist service or GP practice to another, a new shared care agreement must be completed. Ensure that the specialist is informed in writing of any changes to the patient's GP or their contact details.		
15. References Back to top		
- eBNF accessed via https://bnf.nice.org.uk/ on 05/10/2021. - Azathioprine 50mg tablets (Imuran®). Aspen. Date of revision of the text: April 2021. Accessed via https://www.medicines.org.uk/emc/product/3823 on 08/07/2021. - Azathioprine 50mg tablets (Azapress®). Ennogen. Date of revision of the text: 29/06/2016. Accessed via www.medicines.org.uk/emc on 01/10/2020.		

- Mercaptopurine 50mg tablets. Aspen. Date of revision of the text: August 2019. Accessed via <https://www.medicines.org.uk/emc/product/4655/smpc> on 06/10/2021.
- Mercaptopurine 20mg/ml oral suspension (Xaluprine®). Nova Laboratories Ltd. Date of revision of the text: January 2021. Accessed via <https://www.medicines.org.uk/emc/product/2949/smpc> on 06/10/2021.
- British Society of Rheumatology and British Health Professionals in Rheumatology. 2017. [Guidelines for the prescription and monitoring of non-biologic disease-modifying anti-rheumatic drugs](#). Accessed 01/10/2020.
- British Society of Rheumatology and British Health Professionals in Rheumatology. 2016. [Guideline on prescribing drugs in pregnancy and breastfeeding – Part I: standard and biologic disease modifying anti-rheumatic drugs and corticosteroids](#). Accessed 01/10/2020.
- British Association of Dermatologists. 2011. [Guidelines for the safe and effective prescribing of azathioprine](#) (update in progress). Accessed 01/10/2020.
- British Association of Dermatologists. 2007 (updated 2016). [Patient information leaflet: azathioprine](#). Accessed 01/10/2020.
- British Thoracic Society in collaboration with the Thoracic Society of Australia and New Zealand and the Irish Thoracic Society. 2008. [Interstitial lung disease guideline](#) (archived). Accessed 01/10/2020.
- British Society of Gastroenterology. 2019. [Consensus guidelines on the management of inflammatory bowel disease in adults](#). Accessed 01/10/2020.
- British Society of Gastroenterology. 2011. [Guidelines on the management of autoimmune hepatitis](#). Accessed 01/10/2020.
- Association of British Neurologists. 2015. [Myasthenia gravis: management guidelines](#). Accessed 01/10/2020.
- Public Health England. [Immunisation against infectious disease: The Green Book](#). Accessed 01/10/2020.
- Specialist Pharmacy Service. Lactation Safety Information: azathioprine. Reviewed September 2020. Accessed via <https://www.sps.nhs.uk/medicines/> on 08/07/2021.
- Specialist Pharmacy Service. Lactation Safety Information: mercaptopurine. Reviewed September 2020. Accessed via <https://www.sps.nhs.uk/medicines/mercaptopurine/> on 06/10/2021.
- UK Teratology Information Service. Toxbase monograph: Azathioprine or mercaptopurine in pregnancy. December 2019. Accessed via www.toxbase.org on 06/10/2021.
- UK Teratology Information Service. Toxbase monograph: Paternal use of azathioprine or mercaptopurine. December 2019. Accessed via www.toxbase.org on 06/10/2021.
- The Renal Drug Database. Mercaptopurine monograph. Accessed via: <https://renaldrugdatabase.com/monographs/mercaptopurine> on 09/12/21
- NICE Guidance: Crohn's disease: management (NG129) May 2019. Accessed via: <https://www.nice.org.uk/guidance/ng129> on 09/12/21
- NICE Guidance: Ulcerative colitis: management (NG130) May 2019. Accessed via: <https://www.nice.org.uk/guidance/ng130> on 09/12/21

16. Other relevant national guidance

[Back to top](#)

- Shared Care for Medicines Guidance – A Standard Approach (RMOC). Available from <https://www.sps.nhs.uk/articles/rmoc-shared-care-guidance/>
- NHSE guidance – Responsibility for prescribing between primary & secondary/tertiary care. Available from <https://www.england.nhs.uk/publication/responsibility-for-prescribing-between-primary-and-secondary-tertiary-care/>
- General Medical Council. Good practice in prescribing and managing medicines and devices. Shared care. Available from <https://www.gmc-uk.org/ethical-guidance/ethical-guidance-for-doctors/good-practice-in-prescribing-and-managing-medicines-and-devices/shared-care>
- NICE NG197: Shared decision making. Last updated June 2021. <https://www.nice.org.uk/guidance/ng197/>.

17. Local arrangements for referral

[Back to top](#)

Define the referral procedure from hospital to primary care prescriber & route of return should the patient's condition change.

To be agreed and completed locally

- Please note that ESNEFT-Gastroenterology teams on both sites will transfer patients between week 8 to 12 in line with the locally commissioned pathway and monitoring required in the service.

SNEE Integrated Medicines Optimisation Committee approved 14/11/2023 ESNEFT MGG approved 22/11/2023
D&Tc 11/24

Templates for response to shared care guidelines are available in SNEE on DXS and Ardens. The listings are:

Form Name - Appendix 2 Accepting Shared Care

Form Reference - SUF2133

Folder Location - Medicines Management – Care Pathway

Form Name – SNEE Cannot Prescribe Form

Form Reference - SUF2132

Folder Location - Medicines Management – Care Pathway

Appendix 1: Shared Care Request letter (Specialist to Primary Care Prescriber)

Dear [insert Primary Care Prescriber's name]

Patient name: [insert patient's name]

Date of birth: [insert date of birth]

NHS Number: [insert NHS Number]

Diagnosis: [insert diagnosis]

As per the agreed SNEE ICB shared care protocol: [insert shared care guideline reference], this patient is now suitable for prescribing to move to primary care.

The patient fulfils the criteria for shared care, and I am therefore requesting your agreement to participate in shared care. I have completed all the specialist responsibilities listed in the shared care protocol including baseline investigations.

I confirm that his patient has been stabilised on treatment for the agreed amount of time as specified in the shared care guideline

I confirm that I have explained to the patient the risks and benefits of treatment, the rationale and results of the baseline test conducted, the requirements for monitoring, how monitoring will be arranged and the roles of the hospital specialist team, primary care clinicians and the patient in shared care.

Medication name:	
Dose to be continued:	
Date medication started:	
GP to undertake prescribing and monitoring from: (must be at least 1 month after initiation)	
Next blood test due on (if applicable):	

Please respond to this request for shared care, in writing, within 14 days of the request being made where possible.

Kind regards,

[Named Specialist]

Appendix 2: Shared Care Agreement Letter (Primary Care Prescriber to Specialist)

Primary Care Prescriber Response

Dear [insert Doctor's name]

Patient [insert Patient's name]

NHS Number [insert NHS Number]

Identifier [insert patient's date of birth and/or address]

Thank you for your request for me to accept prescribing responsibility for this patient under a shared care agreement and to provide the following treatment

Medicine	Route	Dose & frequency

I can confirm that I am willing to take on this responsibility from *[insert date]* and will complete the monitoring as set out in the shared care protocol for this medicine/condition.

Primary Care Prescriber signature: _____ Date: _____

Primary Care Prescriber address/practice stamp

Appendix 3: Shared Care Refusal Letter (Primary Care Prescriber to Specialist)

Re:

Patient *[insert Patient's name]*

NHS Number *[insert NHS Number]*

Identifier *[insert patient's date of birth and/or address]*

Thank you for your request for me to accept prescribing responsibility for this patient.

In the interest of patient safety NHS *[insert CCG name]*, in conjunction with local acute trusts have classified *[insert medicine name]* as a Shared Care drug, and requires a number of conditions to be met before transfer can be made to primary care.

I regret to inform you that in this instance I am unable to take on responsibility due to the following:

		Tick which apply
1.	The prescriber does not feel clinically confident in managing this individual patient's condition, and there is a sound clinical basis for refusing to accept shared care As the patients primary care prescriber I do not feel clinically confident to manage this patient's condition because <i>[insert reason]</i>. I have consulted with other primary care prescribers in my practice who support my decision. This is not an issue which would be resolved through adequate and appropriate training of prescribers within my practice. I have discussed my decision with the patient and request that prescribing for this individual remain with you as the specialist, due to the sound clinical basis given above.	
2.	The medicine or condition does not fall within the criteria defining suitability for inclusion in a shared care arrangement As the medicine requested to be prescribed is not included on the national list of shared care drugs as identified by RMOC or is not a locally agreed shared care medicine I am unable to accept clinical responsibility for prescribing this medication at this time. Until this medicine is identified either nationally or locally as requiring shared care the responsibility for providing this patient with their medication remains with you	
3.	A minimum duration of supply by the initiating clinician As the patient has not had the minimum supply of medication to be provided by the initiating specialist I am unable to take clinical responsibility for prescribing this medication at this time. Therefore can you please contact the patient as soon as possible in order to provide them with the medication that you have recommended. Until the patient has had the appropriate length of supply the responsibility for providing the patient with their medication remains with you.	

4.	Initiation and optimisation by the initiating specialist As the patient has not been optimised on this medication I am unable to take clinical responsibility for prescribing this medication at this time. Therefore can you please contact the patient as soon as possible in order to provide them with the medication that you have recommended. <i>Until the patient is optimised on this medication the responsibility for providing the patient with their medication remains with you.</i>	
5.	Shared Care Protocol not received As legal responsibility for clinical care lies with the clinician who signs the prescription, I need to ensure that I am in possession of sufficient clinical information for me to be confident to prescribe this treatment for my patient and it is clear where each of our responsibilities lie to ensure the patient is safely managed. For this reason I am unable to take clinical responsibility for prescribing this medication at this time, therefore would you please contact the patient as soon as possible in order to provide them with the medication that you have recommended. <i>Until I receive the appropriate SCP, responsibility for providing the patient with their medication remains with you.</i>	
6.	Other (Primary Care Prescriber to complete if there are other reasons why shared care cannot be accepted)	

I would be willing to consider prescribing for this patient once the above criteria have been met for this treatment.

NHS England 'Responsibility for prescribing between Primary & Secondary/Tertiary care' guidance (2018) states that "when decisions are made to transfer clinical and prescribing responsibility for a patient between care settings, it is of the utmost importance that the GP feels clinically competent to prescribe the necessary medicines. It is therefore essential that a transfer involving medicines with which GPs would not normally be familiar should not take place without full local agreement, and the dissemination of sufficient, up-to-date information to individual GPs." In this case we would also see the term GP being interchangeable with the term Primary Care Prescriber.

Please do not hesitate to contact me if you wish to discuss any aspect of my letter in more detail and I hope to receive more information regarding this shared care agreement as soon as possible

Yours sincerely

Primary Care Prescriber signature: _____

Date: _____

Primary Care Prescriber address/practice stamp